


NASA/TP-2000-210534



Proton-Nucleus Total Cross Sections in Coupled-Channel Approach

*R. K. Tripathi and John W. Wilson
Langley Research Center, Hampton, Virginia*

*Francis A. Cucinotta
Lyndon B. Johnson Space Center, Houston, Texas*

October 2000

The NASA STI Program Office . . . in Profile

Since its founding, NASA has been dedicated to the advancement of aeronautics and space science. The NASA Scientific and Technical Information (STI) Program Office plays a key part in helping NASA maintain this important role.

The NASA STI Program Office is operated by Langley Research Center, the lead center for NASA's scientific and technical information. The NASA STI Program Office provides access to the NASA STI Database, the largest collection of aeronautical and space science STI in the world. The Program Office is also NASA's institutional mechanism for disseminating the results of its research and development activities. These results are published by NASA in the NASA STI Report Series, which includes the following report types:


- **TECHNICAL PUBLICATION.** Reports of completed research or a major significant phase of research that present the results of NASA programs and include extensive data or theoretical analysis. Includes compilations of significant scientific and technical data and information deemed to be of continuing reference value. NASA counterpart of peer-reviewed formal professional papers, but having less stringent limitations on manuscript length and extent of graphic presentations.
- **TECHNICAL MEMORANDUM.** Scientific and technical findings that are preliminary or of specialized interest, e.g., quick release reports, working papers, and bibliographies that contain minimal annotation. Does not contain extensive analysis.
- **CONTRACTOR REPORT.** Scientific and technical findings by NASA-sponsored contractors and grantees.
- **CONFERENCE PUBLICATION.** Collected papers from scientific and technical conferences, symposia, seminars, or other meetings sponsored or co-sponsored by NASA.
- **SPECIAL PUBLICATION.** Scientific, technical, or historical information from NASA programs, projects, and missions, often concerned with subjects having substantial public interest.
- **TECHNICAL TRANSLATION.** English-language translations of foreign scientific and technical material pertinent to NASA's mission.

Specialized services that complement the STI Program Office's diverse offerings include creating custom thesauri, building customized databases, organizing and publishing research results . . . even providing videos.

For more information about the NASA STI Program Office, see the following:

- Access the NASA STI Program Home Page at <http://www.sti.nasa.gov>
- Email your question via the Internet to help@sti.nasa.gov
- Fax your question to the NASA STI Help Desk at (301) 621-0134
- Telephone the NASA STI Help Desk at (301) 621-0390
- Write to:
NASA STI Help Desk
NASA Center for AeroSpace Information
7121 Standard Drive
Hanover, MD 21076-1320

NASA/TP-2000-210534



Proton-Nucleus Total Cross Sections in Coupled-Channel Approach

*R. K. Tripathi and John W. Wilson
Langley Research Center, Hampton, Virginia*

*Francis A. Cucinotta
Lyndon B. Johnson Space Center, Houston, Texas*

National Aeronautics and
Space Administration

Langley Research Center
Hampton, Virginia 23681-2199

October 2000

Available from:

NASA Center for AeroSpace Information (CASI)
7121 Standard Drive
Hanover, MD 21076-1320
(301) 621-0390

National Technical Information Service (NTIS)
5285 Port Royal Road
Springfield, VA 22161-2171
(703) 605-6000

Abstract

Recently, nucleon-nucleon (N-N) cross sections in the medium have been extracted directly from experiment. The in-medium N-N cross sections form the basic ingredients of several heavy-ion scattering approaches including the coupled-channel approach developed at the Langley Research Center. In the present study the ratio of the real to the imaginary part of the two-body scattering amplitude in the medium was investigated. These ratios are used in combination with the in-medium N-N cross sections to calculate total proton-nucleus cross sections. The agreement is excellent with the available experimental data. These cross sections are needed for the radiation risk assessment of space missions.

Introduction

The transportation of energetic ions in bulk matter is of direct interest in several areas including shielding against ions originating from either space radiations or terrestrial accelerators, cosmic ray propagation studies in galactic medium, or radiobiological effects resulting from the work place or clinical exposures. For carcinogenesis, terrestrial radiation therapy, and radiobiological research, knowledge of beam composition and interactions is necessary to properly evaluate the effects on human and animal tissues. For the proper assessment of radiation exposures both reliable transport codes and accurate input parameters are needed. One such important input is total cross sections. The motivation of the work is to develop a method for calculating accurate cross sections. These cross sections are needed in transport methods both deterministic and Monte Carlo.

Nucleon-nucleon (N-N) cross sections are the basic ingredients of many approaches (refs. 1 to 10) to heavy ion scattering problem. Most of the information about these N-N cross sections comes from the free two-body scattering. These cross sections are significantly modified in a nucleus, due to the presence of other nucleons, which is affected through the Pauli exclusion principle and modification of meson field coupling constants. (See ref. 11.) Our theoretical approach is based on the coupled channel method used at the Langley Research Center. (See refs. 1 to 6.) This method solves the Schrödinger equation in eikonal approximation. The method needs modifications at low and medium energies. In earlier work (refs. 12 and 13), the renormalization of the free N-N cross sections were found to be significant at lower and medium energies. These modified in-medium N-N cross sections were used to calculate the total reaction cross sections. The blend of the renormalized N-N cross sections and the coupled-channel method were found to give a reliable basic approach to total reaction cross sections. The purposes of the current paper are as follows:

- (1) To investigate the modification of the ratio of the real to the imaginary part of the two-body amplitude in the medium
- (2) To use these modified two-body amplitudes to calculate total cross sections for proton-nucleus collisions
- (3) To validate and compare the calculated results with the available experimental data
- (4) To provide theoretical results where data are not available because of nonexistence of experimental facilities and/or difficulty in experimental data analysis

Method

The essentials of the coupled-channel method are briefly sketched for completeness. (See refs. 1 to 6 for details.) In this approach, the matrix for elastic scattering amplitude is given by

$$f(\mathbf{q}) = -\frac{ik}{2\pi} \int \exp(-i\mathbf{q}\cdot\mathbf{b}) \{ \exp[i\chi(\mathbf{b})] - 1 \} d^2\mathbf{b} \quad (1)$$

where

f and χ matrices

k projectile momentum relative to center of mass

\mathbf{b} projectile impact parameter vector

\mathbf{q} momentum transfer

$\chi(\mathbf{b})$ eikonal phase matrix

The total cross section σ_{tot} is found from the elastic scattering amplitude by using the optical theorem as follows:

$$\sigma_{\text{tot}} = \frac{4\pi}{k} \text{Im} [f(q=0)] \quad (2)$$

Equations (1) and (2) give

$$\sigma_{\text{tot}} = 4\pi \int_0^\infty b db \{ 1 - e^{-\text{Im}(\chi)} \cos [\text{Re}(\chi)] \} \quad (3)$$

The eikonal phase matrix χ (see refs. 1 to 6 for details) is given by

$$\chi(\mathbf{b}) = \chi_{\text{dir}}(\mathbf{b}) - \chi_{\text{ex}}(\mathbf{b}) \quad (4)$$

The direct and exchange terms are calculated with the following expressions (refs. 1 to 6):

$$\chi_{\text{dir}}(\mathbf{b}) = \frac{APAT}{2\pi k_{NN}} \int d^2\mathbf{q} \exp(i\mathbf{q}\cdot\mathbf{b}) F^{(1)}(-\mathbf{q}) G^{(1)}(\mathbf{q}) f_{NN}(\mathbf{q}) \quad (5)$$

$$\begin{aligned} \chi_{\text{ex}}(\mathbf{b}) &= \frac{APAT}{2\pi k_{NN}} \int d^2\mathbf{q} \exp(i\mathbf{q}\cdot\mathbf{b}) F^{(1)}(-\mathbf{q}) G^{(1)}(\mathbf{q}) \\ &\times \frac{1}{(2\pi)^2} \int d^2\mathbf{q}' \exp(i\mathbf{q}'\cdot\mathbf{b}) f_{NN}(\mathbf{q}+\mathbf{q}') C(\mathbf{q}') \end{aligned} \quad (6)$$

where

$F^{(1)}$ and $G^{(1)}$ projectile and target ground-state one-body form factors, respectively

k_{NN} relative wave number in two-body center-of-mass system

C correlation function (ref. 6)

A_P and A_T mass numbers of projectile and target nuclei, respectively

The two-body amplitude f_{NN} is parameterized as

$$f_{NN} = \frac{\sigma(\alpha + i)}{4\pi} k_{NN} \exp\left(-\frac{Bq^2}{2}\right) \quad (7)$$

where

σ two-body cross section

B slope parameter

α ratio of real part to imaginary part of forward, two-body amplitude

It is well-known that the absorption cross section depends on the imaginary part of the eikonal phase matrix. This leads us to write the two-body amplitude in the medium $f_{NN,m}$ as

$$f_{NN,m} = f_m f_{NN} \quad (8)$$

where f_{NN} is the free NN amplitude and f_m is the system and energy dependent medium multiplier function. (See refs. 12 and 13.) Then the nucleon-nucleon cross sections in the medium ($\sigma_{NN,m}$) can be written as

$$\sigma_{NN,m} = f_m \sigma_{NN} \quad (9)$$

where σ_{NN} is the nucleon-nucleon cross section in free space and the medium multiplier is given by

$$f_m = 0.1 \exp(-E/12) + \left[1 - \left(\frac{\rho_{av}}{0.14}\right)^{1/3} \exp\left(-\frac{E}{D}\right) \right] \quad (10)$$

where E is the laboratory energy in units of A MeV, D is a parameter in units of MeV, as defined subsequently. The numbers 12 and 0.14 are in units of MeV and fm⁻³, respectively. For $A_T \leq 56$ (mass number for iron ion representing heavy elements considered in our transport phenomena),

$$D = 46.72 + 2.21 A_T - (2.25 \times 10^{-2}) A_T^2 \quad (11)$$

and for $A_T > 57$,

$$D = 100 \text{ MeV} \quad (12)$$

In equation (10), ρ_{av} refers to the average density of the colliding system and is

$$\rho_{av} = \frac{1}{2} (\rho_{AP} + \rho_{AT}) \quad (13)$$

where the density of a nucleus $A_i (i = P, T)$ is calculated in the hard sphere model and is given by

$$\rho_{A_i} = \frac{A_i}{(4\pi/3) r_i^3} \quad (14)$$

where the radius of the nucleus r_i is defined by

$$r_i = 1.29 (r_i)_{rms} \quad (15)$$

The root-mean-square radius $(r_i)_{rms}$ is obtained directly from experiment (ref. 14) after subtraction of the nucleon charge form factor (ref. 2).

From equation (3), note that total cross section depends on real component of eikonal phase matrix and, hence (eqs. (5), (6), and (7)), on the product of $\sigma\alpha$ in two-body amplitude. Since the modification of the cross sections in the medium have been determined and tested thoroughly (refs. 12 and 13), the modification of α , ratio of the real to the imaginary part of the two-body amplitude, is studied in the medium to calculate the total cross sections. Some data for total cross sections are available for a few systems at high energies. Unfortunately, no data are available for total cross sections in the low and medium energy range; there are some data for p + Pb in the 100 A MeV range. Therefore, values of the medium-modified α have been tested for higher energies. At low and medium energies, our theoretical results, which incorporate the in-medium two-body amplitudes, can be validated, if and when experimental data become available.

A best estimate of medium-modified α takes into account the enhancement of the cross sections (ref. 15) and stability and is given by

$$\alpha_m = 3 \exp \left[\frac{-(E - 13A^{1/3})^2}{5000} \right] + \frac{K}{1 + \exp[(10 - E)/75]} \quad (16)$$

where

$$K = 0.35 + 0.65 \exp \left[-\frac{2}{3}(N - Z) \right] \quad (17)$$

with N being the neutron number of the nucleus and Z its charge number.

Equation (3) has also been modified to account for the Coulomb force in the proton-nucleus cross sections. This modification has significant effects at low energies and becomes less important as the

energy increases and practically disappears for energies around 50 A MeV and higher.

For nucleus-nucleus collisions, the Coulomb energy is given by

$$V_B = \frac{1.44 Z_P Z_T}{R} \quad (18)$$

where the constant 1.44 is in units of MeV-fm, Z_P and Z_T are charge numbers for the projectile and target, respectively, and R , the radial distance between their centers, is given by

$$R = r_P + r_T + 1.2 \frac{A_P^{1/3} + A_T^{1/3}}{E_{CM}^{1/3}} \quad (19)$$

The number 1.2 in equation (19) is in units of fm-MeV^{1/3}. In our earlier work (refs. 12 and 13), these expressions were used also for the proton-nucleus collisions in order to have a unified picture of any colliding system. However, as shown in the references, equation (19) overestimates the radial distance between proton-nucleus collisions, and hence, equation (18) underestimates the Coulomb energy between them. To compensate for this, we multiplied equation (18) by the following factor (refs. 12 and 13), which gives the Coulomb multiplier to equation (3)

$$X_{Coul} = \left(1 + \frac{C_1}{E_{cm}}\right) \left(1 - \frac{C_2 V_B}{E_{cm}}\right) \quad (20)$$

For $A_T \leq 56$ (mass number for iron),

$$\begin{aligned} C_1 &= 6.81 - 0.17 A_T + (1.88 \times 10^{-3}) A_T^2 \\ C_2 &= 6.57 - 0.30 A_T + (3.6 \times 10^{-3}) A_T^2 \end{aligned} \quad (21)$$

The constant C_1 is in units of MeV. For $A_T > 57$,

$$\begin{aligned} C_1 &= 3.0 \text{ MeV} \\ C_2 &= 0.8 \end{aligned} \quad (22)$$

For the nucleus-nucleus collisions,

$$\begin{aligned} C_1 &= 0 \text{ MeV} \\ C_2 &= 1 \end{aligned}$$

This form of Coulomb energy was found to work well for the proton-nucleus absorption cross sections (ref. 12). Equation (3) is the main equation and is multiplied by equation (20) to get the total cross sections shown in figures 1 to 6.

Results and Conclusions

Figures 1 to 6 show the results of our calculations for the total cross sections for proton on beryllium, carbon, aluminum, iron, lead, and uranium targets, respectively. The experimental data have been taken from the compilation of references 16 and 17. There is paucity of data at lower and intermediate energies (there are some data for $p + \text{Pb}$ in the 100 A MeV range), where the medium modifications play a significant role. For the energy ranges considered, where the data are unavailable, our results provide good theoretical values of total cross sections, since many renormalization effects due to medium, which play an important role in cross sections, have been incorporated in the formalism.

Very good agreement with the experimental results is found for all the systems at higher energies where some data are available. We note that the in-medium cross sections derived earlier in combination with the modified ratio of the real to the imaginary part of the amplitude provide good results for the proton-nucleus total cross sections. It is gratifying to note that the present method gives a consistent basic approach for the total reaction and the total cross sections for the entire energy range for all the systems studied here.

The in-medium two-body amplitudes developed in our approach can be used with ease in other nuclear processes as well.

References

1. Wilson, John W.: Composite Particle Reaction Theory. Ph.D. Diss., College of William and Mary in Virginia, 1975.
2. Wilson, John W.; Townsend, Lawrence W.; Schimmerling, Walter; Khandelwal, Govind S.; Khan, Ferdous; Nealy, John E.; Cucinotta, Francis A.; Simonsen, Lisa C.; Shinn, Judy L.; and Norbury, John W.: *Transport Methods and Interactions for Space Radiations*. NASA RP-1257, 1991.
3. Wilson, John W.; and Costner, Christopher M.: *Nucleon and Heavy-Ion Total and Absorption Cross Section for Selected Nuclei*. NASA TN D-8107, 1975.
4. Cucinotta, Francis A.: Theory of Alpha-Nucleus Collisions at High Energies. Ph.D. Thesis, Old Dominion Univ., 1988.
5. Cucinotta, Francis A.; Townsend, Lawrence W.; and Wilson, John W.: *Target Correlation Effects on Neutron-Nucleus Total, Absorption, and Abrasion Cross Sections*. NASA TM-4314, 1991.
6. Townsend, Lawrence W.: *Harmonic Well Matter Densities and Pauli Correlation Effects in Heavy-Ion Collisions*. NASA TP-2003, 1982.
7. Glauber, R. J.; and Matthiae, G.: High-Energy Scattering of Protons by Nuclei. *Nucl. Phys.*, vol. B21, no. 1, 1970, pp. 135–157.
8. Dadić, I.; Martinis, M.; and Pisk, K.: Inelastic Processes and Backward Scattering in a Model of Multiple Scattering. *Ann. Phys.*, vol. 64, no. 2, 1971, pp. 647–671.
9. Hüfner, J.; Schäfer, K.; and Schürmann, B.: Abrasion-Ablation in Reactions Between Relativistic Heavy Ions. *Phys. Rev. C*, vol. 12, no. 6, 1975, pp. 1888–1898.
10. Feshbach, H.; and Hüfner, J.: On Scattering by Nuclei at High Energies. *Ann. Phys.*, vol. 56, no. 1, 1970, pp. 268–294.
11. Tripathi, Ram K.; Faessler, Amand; and MacKellar, Alan D.: Self-Consistent Treatment of the Pauli Operator in the Brueckner-Hartree-Fock Approach. *Phys. Rev. C*, vol. 8, no. 2, 1973, pp. 129–134.
12. Tripathi, R. K.; Wilson, John W.; and Cucinotta, Francis A.: Nuclear Absorption Cross Sections Using Medium Modified Nucleon-Nucleon Amplitudes. *Nucl. Instrum. & Methods Phys. Res. B*, vol. 145, no. 3, 1998, pp. 277–282.
13. Tripathi, R. K.; Cucinotta, Francis A.; and Wilson, John W.: *Extraction of In-Medium Nucleon-Nucleon Amplitude From Experiment*. NASA/TP-1998-208438, 1998.
14. De Vries, H.; De Jager, C. W.; and De Vries, C.: Nuclear Charge-Density-Distribution Parameters From Elastic Electron Scattering. *At. Data & Nucl. Data Tables*, vol. 36, no. 3, 1987, pp. 495–536.
15. Peterson, J. M.: Nuclear Giant Resonances—Nuclear Ramsauer Effect. *Phys. Rev.*, vol. 125, no. 3, 1962, pp. 955–963.
16. Bauhoff, W.: Tables of Reaction and Total Cross Sections for Proton-Nucleus Scattering Below 1 GeV. *At. Data & Nucl. Data Tables*, vol. 35, 1986, pp. 429–447.
17. Barashenkov, V. S.; Gudima, K. K.; and Toneev, V. D.: Cross Sections for Fast Particles and Atomic Nuclei. *Prog. Phys.*, vol. 17, no. 10, 1969, pp. 683–725.




Figure 1. Total cross sections for proton-beryllium collision as function of energy.




Figure 2. Total cross sections for proton-carbon collision as function of energy.




Figure 3. Total cross sections for proton-aluminum collision as function of energy.




Figure 4. Total cross sections for proton-iron collision as function of energy.




Figure 5. Total cross sections for proton-lead collision as function of energy.




Figure 6. Total cross sections for proton-uranium collision as function of energy.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE October 2000	3. REPORT TYPE AND DATES COVERED Technical Publication	
4. TITLE AND SUBTITLE Proton-Nucleus Total Cross Sections in Coupled-Channel Approach		5. FUNDING NUMBERS WU 101-21-23-03	
6. AUTHOR(S) R. K. Tripathi, John W. Wilson, and Francis A. Cucinotta			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) NASA Langley Research Center Hampton, VA 23681-2199		8. PERFORMING ORGANIZATION REPORT NUMBER L-17994	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Washington, DC 20546-0001		10. SPONSORING/MONITORING AGENCY REPORT NUMBER NASA/TP-2000-210534	
11. SUPPLEMENTARY NOTES Tripathi and Wilson: Langley Research Center, Hampton, VA; Cucinotta: Lyndon B. Johnson Space Center, Houston, TX.			
12a. DISTRIBUTION/AVAILABILITY STATEMENT Unclassified-Unlimited Subject Category 93 Distribution: Standard Availability: NASA CASI (301) 621-0390		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Recently, nucleon-nucleon (N-N) cross sections in the medium have been extracted directly from experiment. The in-medium N-N cross sections form the basic ingredients of several heavy-ion scattering approaches including the coupled-channel approach developed at the Langley Research Center. In the present study the ratio of the real to the imaginary part of the two-body scattering amplitude in the medium was investigated. These ratios are used in combination with the in-medium N-N cross sections to calculate total proton-nucleus cross sections. The agreement is excellent with the available experimental data. These cross sections are needed for the radiation risk assessment of space missions.			
14. SUBJECT TERMS Cross sections; In-medium modifications; Scattering; Heavy ion collision			15. NUMBER OF PAGES 15
			16. PRICE CODE A03
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL